



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

[Jani, Rati](#), Mihrshahi, Seema, Mandalika, Subhadra, & [Mallan, Kimberley M.](#)

(2014)

Validity of maternal-reported weight and height of children 2-5 year olds.
Indian Journal of Pediatrics.

This file was downloaded from: <https://eprints.qut.edu.au/68090/>

© Copyright 2014 Please consult the authors

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<https://doi.org/10.1007/s12098-014-1400-0>

Validity of maternal-reported weight and height of children 2-5 year olds

R Jani, *S Miharshahi, [†]S Mandalika and [‡]KM Mallan

PhD candidate at the Institute of Health & Biomedical Innovation, Queensland University of Technology (QUT), Brisbane, Australia; *Lecturer at the School of Population Health, The University of Queensland, Brisbane, Australia, [†]Assistant professor at the College of Home Science, Nirmala Niketan, affiliated to University of Mumbai, India, [‡]Senior research fellow at the Institute of Health & Biomedical Innovation and School of Exercise and Nutrition Sciences, QUT, Brisbane, Australia

To the Editor: In affluent-urban areas of India, overweight (6%) and obesity (8%) is prevalent in children as young as 2-5 years [1]. A potential risk factor for childhood obesity could be parent's under-reporting their child's anthropometry. In Indian culture a larger body size is typically acceptable, and mothers may consider a chubby baby as healthy [2]. Therefore, it was proposed that Indian mothers may under-report their child's weight status. The present study examined the validity of maternal reported height and weight of young, urban-affluent Indian children aged 2-5 years.

After receiving approval from the QUT Human Research Ethics Committee, Australia 111 mothers with children aged 2-5 years attending private medical clinics (n=5) in the affluent areas of Mumbai were recruited. Child height and weight were measured by the researcher using standard equipment/protocols. Mothers also reported their child's height and weight.

Mothers significantly under-reported their child's height (mean difference: -11.3 ± 22.2 cm, $t(83) = -4.7$, $p < 0.001$). No difference was observed between measured and reported mean weight (mean difference: 0.0 ± 3.9 kg, $p = 0.91$). Only the weight data was further investigated by categorizing the child's weight status according to Weight-for-Age Z-scores cut-offs [3]. There

was no difference between measured and maternal reports for underweight (32.4 % vs 30.0 %, $p=0.89$) and healthy weight (59.5 % vs 69.1 %, $p=0.25$) children. In contrast, a lower proportion of children were classified as overweight/obese based on maternal reports (8.1 % vs 0.9 %, $p=0.02$). These findings were further confirmed as both the sensitivity and specificity for identifying overweight/obesity in children were significantly lower (90.0 % vs 10.0 %, $p=0.02$ and 51.7 % vs 48.3 %, $p=0.02$, respectively) using maternal reported weight compared with measured weight.

This study showed that mothers under-reported their child's height and overweight/obese status. The results imply that parents who underreport their child's overweight/obese status may have poor awareness about their child's true weight, and thus may be less willing to adhere to health professionals' weight management recommendations and participate in interventions promoting a healthy lifestyle [4]. In conclusion, this study confirms the importance of using measured height and weight data in this population.

Funding: The study was a part of R Jani's doctoral project funded by QUT, Australia. Dr. KM Mallan occupied the Heinz Postdoctoral Research Fellowship.

Author's contributions: All authors were involved in each phase of this study including data collection, analysis and interpretation, and preparation of the manuscript. All authors gave their final approval for submission.

Declaration: The authors declare that they have no conflict of interests.

Acknowledgements: The study was a part of R Jani's doctoral project funded by QUT, Australia. Dr. KM Mallan occupied the Heinz Postdoctoral Research Fellowship. We would like to acknowledge the contribution to this project by Prof. Lynne Daniels from School of Exercise & Nutrition Sciences, QUT, Australia. All authors supervised and gave input during data collection, recruitment and preparation of the manuscript and provided their final approval for submission. The authors have no conflicts of interest.

References:

1. Khadilkar V, Khadilkar A, Cole T, Chiplonkar S, Pandit D. Overweight and obesity prevalence and body mass index trends in Indian children. *Int J Pediatr Obes.* 2012; 6:e216-e25.
2. Bhave S, Bavdekar A, Otiv M. IAP National task force for childhood prevention of adult diseases: childhood obesity. *Indian Pediatr.* 2004; 41:559-75.
3. World Health Organization. Child growth standards: training course on child growth assessment. WHO, Geneva; 2008.
4. Parry A, Netuveli G, Parry J, Saxena S. A systematic review of parental perception of overweight status in children. *J Ambul Care Manage.* 2008; 3:253-68.